

Lifeline's Helping someone at risk of suicide tool kit provides information about the following:

Are you concerned that someone close to you is considering taking their own life?

Have you noticed changes in their attitude and behaviour?

Has someone you know attempted suicide?

Would you like to know how to help them stay safe and seek help?

It is distressing to realise that someone close to you may be considering suicide. This tool kit will help you identify signs to look for, decide what to do and where you can go for help.

Most people who consider suicide get through the crisis. Family, friends and professionals can make a big difference in helping people stay safe and re-establish reasons for living.

Are you thinking of suicide?


You are not alone. Thoughts of suicide occur for many people for a range of reasons. The most important thing to remember is that help is available. Talking to someone is a good place to start, even though it may seem difficult. Approach someone you trust or call one of the helpline numbers listed at the end of this tool kit. Tell someone today if you are thinking about suicide.

Why does someone consider suicide?

People considering suicide often feel very isolated and alone. They may feel that nobody can help them or understand them. They believe that suicide is the only way out of the difficulties that they are experiencing.

Having a mental health problem does not mean a person will have thoughts of suicide – many don't. However, mental health problems can affect the way people view situations. They affect motivation and openness to seek help, therefore we need to be particularly aware of the vulnerability to suicide that people experiencing mental illness can have.





What do I do now?

The following tips will help you know what to do

1

Do something now:

If you are concerned that someone you know is considering suicide, act immediately. Don't assume that they will get better without help or that they will seek help on their own. It's easy to avoid being part of that help or to hope that someone else will step in. Reaching out now could save a life.

2

Acknowledge your reaction:

When you realise that you need to take action to help someone who is considering suicide, your natural reaction may be to:

- Panic
- Ignore the situation and hope it will go away
- Look for quick-fix solutions to make the person feel better
- Criticise or blame the person for their feelings
- Tell the person they are being silly and trivialise the issue or dismiss them

These reactions are common but not helpful. It's natural to feel panic and shock, but take time to listen and think before you act. If you find you're really struggling, enlist the help of a trusted friend or helpline.

3 Be there for them:

Spend time with the person and express your care and concern. Ask them how they are feeling, hear their pain and listen to what's on their mind. Let them do most of the talking.

4 Ask if they are thinking of suicide:

Unless someone tells you, the only way to know if a person is thinking of suicide is to ask. Asking can sometimes be very hard but it shows that you have noticed things, been listening, that you care and that they're not alone. Talking about suicide will not put the idea into their head but will encourage them to talk about their feelings. They will often feel a great sense of relief that someone is prepared to talk with them about their darkest thoughts.

5 Check out their safety:

If a person is considering suicide it is important to know how much they have thought about it, so ask them about the following:

- Have they thought about how and when they plan to kill themselves?
- Are they able to carry out their plan?
- Have they ever deliberately harmed themselves?
- What support can they access to stay safe and get help?
- How can you help them draw on connections with family, friends, pets, religious convictions, personal coping strengths and strategies?

Use this information to decide what to do. If you are really worried, don't leave the person alone. Seek immediate help – see contact numbers listed on Lifeline's website www.lifeline.org.au or phone Lifeline on 13 11 14.

Remove any means of suicide available, including weapons, medications, alcohol and other drugs, even access to a car. Be aware of your own safety.

For immediate crisis intervention when life may be in danger ring the ambulance or police on 000 or go to your local hospital emergency department.

6 Decide what to do:

Now that you have this information you need to discuss together what steps you are going to take. What you decide to do needs to take into account the safety concerns that you have. Don't agree to keep it a secret.

You may need the help of others to persuade the person to get professional help – or at least take the first steps to stay safe. These may include their partner, parents, or close friends. Only by sharing this information can you make sure that the person gets the help and support they need.

Sometimes the person at risk says they don't want help. Yet we know most people are in two minds about suicide. Make keeping them safe your first priority. Consider the long-term benefits of getting help for the person. It may mean risking the relationship you have with them, but you could be saving a life.

7 Take action:

The person can get help from a range of professional and supportive people:

- GP
- Counsellor, psychologist, social worker
- School counsellor, youth group leader, sports coach
- Emergency services – police and ambulance
- Mental health services
- Community health centres
- Priest, minister, religious leader
- Telephone crisis support services such as Lifeline and Kids Helpline

When the person has decided who they are most willing to tell, help them prepare what they will say. Many people find it difficult to express their suicidal thoughts.

Offer to accompany the person to the appointment. After the appointment, check that they raised the issue of suicide and ask what help they were offered. Help them follow through with the recommendations.

In some situations the person may refuse to get help. While it's important that you find them the help they need, you can't force them to accept it. You need to ensure that the appropriate people are aware of the situation. Don't shoulder this responsibility alone.

8

Ask for a promise:

Thoughts of suicide often return and when they do it is important for the person to again reach out and tell someone. Asking them to promise to do this makes it more likely that it will happen. Encourage the person to promise to call you, a GP or Lifeline on 13 11 14 if suicidal thoughts return, and to do this before they harm themselves.

9

Look after yourself:

If you're helping someone who is considering suicide, make sure you also take care of yourself. It is difficult and emotionally draining to support someone who is suicidal, especially over an extended period.

- Don't do it on your own. Find someone to talk to - friends, family or a professional.
- Recruit other people to help support the person you are worried about.
- Get in touch with carer organisations or support groups. Contact Lifeline on 13 11 14 to find what's available in your area.
- Try not to let your concerns about the other person dominate your life. Make sure you continue to enjoy your usual activities, take time out to have fun and keep a sense of perspective.

10

Stay involved:

Thoughts of suicide don't easily disappear without the person at risk experiencing some change. Their situation or feelings may change, or they may feel more supported and able to deal with it. In either situation, the continued involvement of family and friends is very important. Below are some tips to ensure the person at risk continues to get the best help possible:

- Ensure the person has 24 hour access to some form of support. This may be you, other family members and friends, or Lifeline.
- Accompany the person to appointments if possible. Your support can be a great encouragement.
- If you are the primary carer, try to establish a good relationship with the health professionals responsible for the person's treatment. Your opinion and input is valid and may be very valuable.
- Advocate for the person. Sometimes a service or health professional may not be capable of meeting all the person's needs. You can advocate for appropriate services.
- Discuss with the person what issues or situations might trigger further suicidal thoughts. Plan how to reduce this stress and what coping strategies can be used.
- Continue to be supportive but not overprotective.
- Encourage the person to write out a plan for how they are going to stay safe, the steps they will take and other people to get involved if things start to get rough. A clearly documented stepped plan is a useful tool to keep a person safe.

Mental Health

People who have recently been discharged from hospital for treatment of mental health problems may also be at higher risk of suicide. It is important that they receive ongoing support in the community.

You may be able to help, by supporting them to attend any follow-up visits with their GP or mental health specialists. If the person has not sought help for a suspected mental health problem, you should encourage them to do so.





What to look out for:

Situations - what's happening in the person's life? Have they experienced any life changes recently?

- Recent loss (a loved one, a job, an income/ livelihood, a relationship, a pet)
- Major disappointment (failed exams, missed job promotions)
- Change in circumstances (separation/divorce, retirement, redundancy, children leaving home)
- Mental disorder or physical illness/injury
- Suicide of a family member, friend or a public figure
- Financial and/or legal problems

Feelings – how does the person feel about their life?

Events and life changes can be difficult and sometimes devastating. Most people who experience them don't consider suicide, but some do. Be aware of:

- How the person feels about what's happened
- What it means to them
- Whether the pain feels bearable

Behaviours – what are they doing?

People at risk of suicide usually give clues by the way they behave. These may include:

- Previous suicide attempts
- Being moody, sad and withdrawn
- Talking of feeling hopeless, helpless or worthless
- Taking less care of themselves and their appearance
- Losing interest in things they previously enjoyed
- Difficulty concentrating and/or sleeping
- Being more irritable or agitated
- Talking or joking about suicide/death
- Expressing thoughts about death through drawings, stories, songs etc.
- Saying goodbye to others and/or giving away possessions
- Engaging in risky or self-destructive behaviour
- Increasing alcohol/drug use

Places to go for help now:

For immediate crisis intervention when life may be in danger call the ambulance or police on 000 or go to your local hospital emergency department.

For further information about places to go for help when someone is at risk of suicide, visit the Lifeline website www.lifeline.org.au

National Helpline Support

- Lifeline 13 11 14
- Kids Helpline 1800 55 1800
- Mensline Australia 1300 78 99 78
- Salvation Army Hope Line for suicide bereavement support 1300 467 354
- The Suicide Call Back Service 1300 659 467

Acknowledgements:



Part of your Community

Prime Super is the proud sponsor of the Lifeline Information Service – your mental health and self-help resource.

Prime Super is Australia's only nationally operating super fund dedicated to rural and regional Australia.

For more information on Prime Super, please ring 1800 675 839 or visit their website www.primesuper.com.au

Lifeline and Prime Super are working in partnership to promote mental health awareness, help-seeking and suicide prevention.



For 24 hour telephone crisis support call 13 11 14
For more information visit www.lifeline.org.au
To donate call 1800 800 768

This Tool Kit has been produced by the Lifeline Information Service as a public service. You are welcome to reproduce it without alteration and with acknowledgement of Lifeline. We invite your feedback and comments at infoservice@lifeline.org.au

Last revised June 2010